

PERITONITIS (Continued).

1018—℞ Magnesii sulphatis, . . . ℥iiss.

In pulv. no. xii div.

Sig.: A powder in hot peppermint-water every hour until the bowels are freely opened. (*In acute peritonitis, at the beginning of the attack.*)

MUNDÉ, *Annual Univ. Med. Sci.*

1019—℞ Morphiæ sulphatis, . . . gr. iv.
Aquæ destillatæ, . . . ℥ij. —M.

Sig.: Ten or fifteen minims as required, hypodermically, to control the vomiting. Give no food for twenty-four or forty-eight hours. (*Especially in peritonitis following surgical operations.*)

TAIT, *Annual Univ. Med. Sci.*

PERTUSSIS (See Whooping-Cough).

PHAGEDENA.

1020—℞ Saloli, gr. v-l.
Amyli, ℥j. —M.

Ft. pulv.

Sig.: Dust over locally.

SEIFERT.

1021—℞ Iodoformi,
Chinoidini,
Ferri redacti, āā ℥j.—M.

In pil. no. xx div.

Sig.: One pill three times daily. (*In sloughing phagedena.*)

BARTHOLOW.

1022—℞ Iodoformi, ℥iiss.
Thymoli, ℥v.
Sacchari lactis, gr. ij.—M.

Ft. pulv.

Sig.: Dust over sores.

HOWARD.

1023—℞ Acidi pyrogallici, ℥ij.
Pulv. amyli, ℥j.—M.

Ft. pulv.

Sig.: Dust over or insufflate the sores twice daily. The powder should be fresh, and kept in a tightly-corked bottle.

TERRILLON.

1024—℞ Acidi nitrici fort., ℥j.

Sig.: Apply thoroughly but carefully to the whole secreting surface, after drying it. Give the patient an anæsthetic if necessary, and a hypodermic of morphia later.

VAN BUREN and KEYES.