

PHTHISIS (Continued).

1040—R Acidi tannici, ℥ij.
Glycerinæ, ℥j.—M.

Sig.: A teaspoonful from two to four times daily.
ARTHAUD and RAYMOND, Annual Univ. Med. Sci.

1041—R Antipyrin., ℥j.
Aquæ ferventis, ℥iij.—M.

Sig.: From thirty minims to four drachms daily by hypodermic, according to strength of the patient.
ZAKRZHEVSKI, Annual Univ. Med. Sci.

1042—R Cupri acetatis, gr. ij.
Sodii carbonatis, gr. xij.
Glycerinæ et ext. glycyrrhizæ, q. s. —M.

Ft. massa et in pil. no. xii div.
Sig.: One pill night and morning on an empty stomach.
LUTON, Annual Univ. Med. Sci.

1043—R Pulv. agarici, ℥j.

In pulv. no. xii div.
Sig.: One powder every two hours (for three doses) if necessary. (*For night sweats.*)
A. PETER, Annual Univ. Med. Sci.

1044—R Ext. ergotæ fld., ℥j.

Sig.: Twenty drops three times daily. (*To relieve diarrhœa and sweats.*)
A. L. HODGSON, Annual Univ. Med. Sci.

1045—R Atropiæ sulphatis, gr. j.
Aquæ destillatæ, ℥v.—M.

Sig.: Twelve minims hypodermically. (*For hæmoptysis; desperate cases.*)
HAUSMANN.

1046—R Acetphenetidin., ℥iiss.

In pulv. no. xv div.
Sig.: A powder stirred in a little water two or three times daily. (*For hectic.*)
HINSBERG and KAST, Annual Univ. Med. Sci.

1047—R Antifebrin., ℥ij.
Spts. vini gallici, ℥iij.—M.

Sig.: From a dessertspoonful to a tablespoonful two or three times daily. (*For hectic of phthisis.*)
FAUST, Annual Univ. Med. Sci.