

PHTHISIS (Continued).

1048—R Ammonii boratis, ʒij.
Syr. simplicis,
Elix. simplicis, āā ʒiiss.—M.

Sig.: A teaspoonful thrice daily. (*For expectoration and hectic.*) LASHKEVITCH, Annual Univ. Med. Sci.

1049—R Olei delphinidæ (porpoise-oil), Oss.

Sig.: A teaspoonful to a tablespoonful thrice daily, after meals. (*Aliment in phthisis.*)
WEST, Annual Univ. Med. Sci.

1050—R Pyridin, ʒj.

Sig.: Six to ten drops, increased to twenty-five drops daily, in three-drop doses. In urgent cases three to five drops may be inhaled directly from a handkerchief. (*For dyspnœa.*) DE RENZI, Annual Univ. Med. Sci.

1051—R Antipyrin, gr. lxxv.
Spts. jamaicensis, ʒv.
Syr. limonis, ʒviiss.
Aquæ destillatæ, ʒiiss.—M.

Sig.: One to three tablespoonfuls once or twice daily. (*For hectic.*)

1052—R Terebene, ʒiv.
Pulv. acaciæ, ʒij.
Syr. zingiberis, ʒviiss.
Aquæ, ʒxv.—M.

Sig.: A teaspoonful three times daily. (*Relieves dyspnœa and flatulence.*) VIGIER.

1053—R Terpene hydratis, gr. lxxv.
Spts. vini rectificati (95 per cent.), ʒv.
Glycerinæ, ʒx.—M.

Sig.: A teaspoonful or two in a little sweetened or aromatized water two or three times daily. (*Expectorant.*) VIGIER.

1054—R Terpinol,
Sodii benzoatis, āā gr. xv.
Sacchari albi, q. s. —M.

In capsulas no. x div.

Sig.: A capsule every hour or two. (*To diminish the expectoration and remove its odor.*) RABOW.