

**PHTHISIS (Continued).**

1055—R Amyl hydratis, . . . . . gr. cv.  
Ext. glycyrrhizæ, . . . . . ℥iiss.  
Aquæ destillatæ, . . . . . ℥xv.—M.

Sig.: The half to be taken at bed-time. (*For the insomnia of phthisis.*) FISCHER.

1056—R Amyl hydratis, . . . . . ℥iiss.  
Morphiæ muriatis, . . . . . gr.  $\frac{1}{3}$ .  
Ext. glycyrrhizæ, . . . . . ℥iiss.  
Aquæ destillatæ, . . . . . ℥xv.—M.

Sig.: The half to be taken at bed-time. (*Insomnia of phthisis.*) FISCHER.

1057—R Creasoti, . . . . . ℥iiss.  
Tinct. gentianæ, . . . . . ℥j.  
Spts. vini rectificati, . . . . . ℥viiij.  
Vini xerici, . . . . . ad Oij.—M.

Sig.: A tablespoonful in a wineglassful of water three times daily. (*In incipient tuberculosis.*) FRÄNTZEL.

1058—R Sodii iodidi, . . . . . gr. lxxv.  
Sodii bromidi, . . . . . ℥iiss.  
Sodii chloridi, . . . . . ℥v.  
Aquæ destillatæ, . . . . . ℥l.—M.

Sig.: A teaspoonful every morning in a cupful of milk. "Summer Cod-liver Oil." Contains the principal constituents of olei morrhuæ. (*Aliment in phthisis.*) POTAIN.

1059—R Spts. vini gallici vel jamai-  
censis, . . . . . ℥iiss.  
Olei menthæ pip., . . . . . ℥j.  
Glycerinæ, . . . . . ℥x.—M.

Sig.: To be taken in divided doses during the day. In cases which present no sign of abnormal excitability of the nervous system or heart, the dose of glycerin may be raised to twelve or fifteen ounces daily. (*Aliment, when patients cannot take olei morrhuæ.*) JACCOUD.

**PILES (See Hemorrhoids).**

**PITYRIASIS (See Skin Diseases).**