

PLEURISY (Continued).

1067—R Acidi tannici, . . . gr. xxx.
Confection. rosæ, . . . q. s. ut ft. massa.

In pil. no. xv div.

Sig.: Two to four pills morning and evening. (*In purulent effusion.*)
DEBOUÉ.

1068—R Potassii iodidi, . . . ℥iv.
Aquæ, . . . ℥vj.—M.

Sig.: One teaspoonful in milk every four hours, with the following:—

1069—R Tinct. iodinii co., . . . ℥iij.

Sig.: Divide the surface of the affected side into three sections, and paint one section each day. (*For chronic pleuritic effusion.*)
BARTHOLOW.

1070—R Magnesii sulphatis, . . . ℥vi-vij.

In pulv. no. viii div.

Sig.: A powder in two tablespoonfuls of water before food, and no fluids for some time afterward. (*In pleuritic effusion.*)
M. HAY.

1071—R Collodii cum cantharide, . . . ℥ss.

Sig.: Apply with a brush over a small area, heal quickly, and repeat. (*In pleuritic effusion.*)

RINGER.

1072—R Syr. ferri iodidi, . . . ℥ij.
Syr. simplicis, . . . ℥ij.—M.

Sig.: A teaspoonful every two hours, with the following:—

1073—R Iodinii, . . . ℥ss.
Potassii iodidi, . . . ℥ij.
Aquæ destillatæ, . . . ℥ij.—M.

Sig.: Apply on the affected side of the chest.

NIEMEYER.

1074—R Pulv. sinapis, . . . ℥ss.
Pulv. lini, . . . ℥viiij.
Aquæ bullientis, . . . q. s.—M.

Ft. cataplasma

Sig.: Make the poultice so wet that it moistens the hands in holding it. Place it between two pieces of muslin, cover with oiled muslin, and renew when beginning to cool. (*In pleurisy of children.*)

J. LEWIS SMITH.