

QUINSY (Continued).

1148—R Sodii salicylatis, ℥iii-iv.
Syr. acaciæ, ℥ss.
Aquæ cinnamomi, ad ℥ij.—M.

Sig.: A dessertspoonful every three hours.
EASBY, *Annual Univ. Med. Sci.*

1149—R Salinaphthol, gr. xx-xxv.
Spts. vini rectificati, ℥j. —M.

Sig.: One part to twenty of water, as an antiseptic gargle.
GEORGI, *Annual Univ. Med. Sci.*

1150—R Sodii bicarbonatis, ℥ss.

Sig.: Apply locally to the tonsil in powder, or in warm solution.
BAKER, *Annual Univ. Med. Sci.*

1151—R Acidi boracici, ℥ij.
Eau de Pagliari, ℥x.
Aquæ, ℥vii ℥viss.

Ft. gargarisma. —M.

Sig.: Use as a gargle several times daily, and each time follow by an application of—

1152—R Iodoformi, ℥j.
Collodii flexilis, ℥vij.—M.

Sig.: "Iodoform collodion."
LEBRUN, *Annual Univ. Med. Sci.*

RACHITIS (RICKETS), SCROFULA, STRUMA.

1153—R Olei morrhuæ, ℥v.
Phosphori, gr. iij.—M.

Ft. sol.

Sig.: One teaspoonful daily.
KASSOWITZ, *Annual Univ. Med. Sci.*

1154—R Phosphori, gr. iij.
Solve in—
Carbonis sulphureti, ℥lxxv.
Aquæ destillatæ, ℥v. —M.

Sig.: Half a teaspoonful twice daily. The bottle must be kept tightly corked. No sugar must be combined with the preparation, but may be given after taking the dose.

HASTERLIK, *Annual Univ. Med. Sci.*