

**RACHITIS (RICKETS), SCROFULA, STRUMA (Continued).**

1171—R Potassii iodidi, . . . . . gr. xxx.  
Tinct. iodinii, . . . . .  
Acidi tannici, . . . . . āā gr. xv.  
Syr. quiniæ, . . . . . ℥viiss.  
Syr. acaciæ, . . . . . ℥iv ℥vss. —M.

Sig.: A fourth part to be taken every two hours until four doses are taken by an adult. For children, a corresponding fractional dose. GUIBOUT.

**RATTLESNAKE-BITE.**

1172—R Potassii permanganatis, . . . . . ℥ss.  
Aquæ destillatæ, . . . . . ℥iij.—M.

Sig.: Apply to the wound, and inject hypodermically above the seat of injury. At the same time take internally the following:—

1173—R Aquæ ammoniæ, . . . . . ℥iv.

Sig.: A half-teaspoonful in water, repeated every ten or fifteen minutes.

HAWACK and ARBOC, Annual Univ. Med. Sci.

1174—R Aquæ ammoniæ, . . . . . ℥j.  
Aquæ, . . . . . ℥iij.—M.

Sig.: Inject thirty minims hypodermically into a superficial vein above seat of injury. HALFORD.

1175—R Hydrargyri chloridi corrosivi, gr. j.  
Potassii iodidi, . . . . . gr. ij.  
Brominii, . . . . . ℥iiss.  
Alcoholi diluti, . . . . . ℥xxx.—M.

Ft. sol.

Sig.: A teaspoonful in wine or brandy as often as necessary. BIBRON.

**REMITTENT FEVER (See Fever).**

**RENAL CALCULI (See Calculi).**

**RENAL DROPSY (See Dropsy).**

**RENAL HEMORRHAGE (See Hæmaturia).**

**RHEUMATISM, ACUTE.**

1176—R Antifebrin., . . . . . ℥j.

Fiant capsulæ no. xvi.

Sig.: One or two capsules three to six times daily.

EISENHART, Annual Univ. Med. Sci.