

RHEUMATISM, ACUTE (Continued).

1177—R Antipyrin., ℥iij.
Syr. aurantii cort., ℥j.
Aquæ, ad ℥iij.—M.

Sig.: A dessertspoonful in water thrice daily. (*In afebrile cases.*)
GERMAIN SÉE.

1178—R Hydrochinon., ℥ss.
Aquæ cinnamomi, ℥iij.—M.

Sig.: One-half to three teaspoonfuls two to four times daily, until the fever abates.
SYLVESTRINI and PICCHINI, *Annual Univ. Med. Sci.*

1179—R Liq. ammonii ichthyosulphatis
(30 per cent.), ℥ij.
Lanolini, ℥j.—M.

Ft. unguentum.
Sig.: To be rubbed over the swollen joints. Take internally the following:—

1180—R Ichthyol., ℥j.

Fiant capsulæ no. xx.
Sig.: Three to six capsules during the twenty-four hours. (*In both acute and chronic cases.*)
SCHMIDT, *Annual Univ. Med. Sci.*

1181—R Acidi salicylici, ℥iij.
Potassii bicarbonatis, ℥vj.
Aquæ, ℥iij.—M.

Sig.: A teaspoonful every three hours.
DONNELLY.

1182—R Pimentæ, ℥vi ℥ij.
Aquæ ammoniæ, ℥iii ℥j.
Ess. thymi,
Chloral hydratis, āā ℥iiss.
Spts. vini rectificati (60°), Oij.—M.

Ft. linimentum.
Sig.: "Apone." Use pure or mixed with olive oil. (*For friction about rheumatic joints.*)
POULET.

1183—R Sodii salicylatis, ℥ss.
Tinct. lavandulæ co., ℥iv.
Glycerinæ, ℥ss.
Aquæ, ad ℥viiij.—M.

Sig.: A tablespoonful every hour or two until pain and fever abate; then at longer intervals.
F. MINOT, *Mass. Genl. Hosp.*