

**RHEUMATISM, ACUTE (Continued).**

1184—R Acidi salicylici, . . . . . ℥ss.  
Ferri pyrophosphatis, . . . . . ℥j.  
Sodii phosphatis, . . . . . ℥x.  
Aquæ, . . . . . ℥vj.—M.

Sig.: A tablespoonful every two hours until the improvement justifies less frequent doses, or unless constitutional effects are produced.

G. L. PEABODY, *New York Hosp.*

1185—R Sodii salicylatis, . . . . . ℥vj.  
Glycerinæ, . . . . . ℥iv.  
Aquæ cinnamomi, . . . . . ad ℥vj.—M.

Sig.: A tablespoonful every two or three hours until tinnitus aurium is produced; then every four to six hours until the acute symptoms have abated. Then give—

1186—R Sodii bicarbonatis, . . . . . ℥iv-vj.

In pulv. no. xii div.

Sig.: A powder in a half-glassful of water every four hours, until the urine is alkaline to test-paper. If the patient is anæmic, omit the salicylate and begin on the soda at once, and give cod-liver oil and iron from the first. (*In robust cases.*)

A. L. LOOMIS, *Bellevue Hospital, N. Y.*

1187—R Olei gaultheriæ, . . . . . ℥j.

Sig.: Fifteen or twenty minims to be given in capsules or floated on milk or water every two hours, until the acute symptoms abate; then gradually diminish to one drachm daily until convalescence; then combine iron. If any joint-stiffness remains, then give—

1188—R Lithii salicylatis, . . . . . ℥ii-ij.

Sig.: To be given, dissolved in water, during the twenty-four hours.

KINNICUTT, *St. Luke's Hospital, N. Y.*

1189—R Sodii salicylatis,  
Potassii citratis, . . . . . āā gr. xv.  
Aquæ, . . . . . ℥ss. —M.

Sig.: To be given every two hours until the pain and fever abate. Also the following:—