

**RHEUMATISM, ACUTE (Continued).**

1190—R Liq. opii sedativi, . . . . . ℥j.  
Potassii bicarbonatis, . . . . . ℥iv.  
Glycerinæ, . . . . . ℥ij.  
Aquæ bullientis, . . . . . ℥ix.—M.

Sig.: "Fuller's Lotion." Soak a piece of flannel or spongiopiline in the above hot solution, and wrap it around the painful joint.

OSLER, *Univ. of Penna., Phila.*

1191—R Acidi salicylici, . . . . . ℥iij.  
Sodii boratis, . . . . . gr. xv.  
Aquæ menthæ pip., . . . . . ad ℥vj. —M.

Ft. sol.

Sig.: One-third to be taken during twenty-four hours. If there be no improvement in three or four days, discontinue and use—

1192—R Ammonii bromidi, . . . . . ℥iii-iv.

In pulv. no. xii div.

Sig.: A powder in a half-glassful of water every four hours. When the acute symptoms abate, add twelve to sixteen grains of quinine daily.

DA COSTA, *Jefferson Hospital, Phila.*

1193—R Propylaminæ, . . . . . gr. xxiv.  
Aquæ menthæ pip., . . . . . ℥vj. —M.

Sig.: A tablespoonful every two or three hours.

JAS. TYSON, *Phila. Hosp.*

1194—R Sodii salicylatis, . . . . . gr. xv.  
Sodii bicarbonatis, . . . . . gr. xxx.  
Aquæ menthæ pip., . . . . . ℥ss. —M.

Sig.: To be taken every third or fourth hour. When the acute symptoms abate, then give—

1195—R Mist. ferri et ammonii acetatis  
(U. S. P.), ℥iv.

Sig.: A dessertspoonful or two in a wineglassful of water, thrice daily.

J. C. WILSON, *Phila. Hosp.*

**RHEUMATISM, CHRONIC.**

1196—R Calcii chloridi, . . . . . ℥iij.  
Syr. simplicis, . . . . . ℥iv.  
Olei gaultheriæ, . . . . . gtt. iv.—M.

Sig.: A tablespoonful thrice daily for adults. One-third dose for children. Also use externally—