

SARCINÆ AND TORULÆ.

1216—R Sodii sulphitis, gr. xxx-xl.
Infusi quassiaë, ℥iiss. —M.

Ft. haustus.

Sig.: To be taken thrice daily. SIR W. JENNER.

1217—R Sodii hyposulphitis, ℥vj.
Infusi quassiaë, ℥xij.—M.

Sig.: A teaspoonful thrice daily. R. NEALE.

1218—R Acidi sulphurosi, ℥ss-j.
Aquaë, ℥ij.—M.

Ft. haustus.

Sig.: To be taken thrice daily. TANNER.

1219—R Acidi sulphurosi, ℥i-iss.
Infusi calumbæ, ℥xij.—M.

Ft. haustus.

Sig.: A wineglassful ten minutes before meals.
LAWSON.

SATYRIASIS (See Nymphomania).

SCABIES (See Lice).

SCARLATINA (See also Fever and Diphtheria).

1220—R Tinct. ferri chloridi, ℥ij.
Potassii chloratis, ℥i-ij.
Syr. simplicis, ℥iv.—M.

Sig.: A teaspoonful every hour or two to a child of
four or five years. J. LEWIS SMITH.

1221—R Ammonii carbonatis, ℥i-iss.
Syr. simplicis, ℥j.
Aquaë, ad ℥iiss.—M.

Sig.: A teaspoonful every hour, or every two or
three hours, according to the severity of the case.

PEART.

1222—R Infusi digitalis, ℥iv.

Sig.: One-half to one teaspoonful every two, three,
or four hours. BARTHOLOW.

1223—R Tinct. aconiti radiceis, ℥ss.

Sig.: One-half to one drop in a teaspoonful of water
every quarter-hour for two hours; afterward hourly.
If there is much prostration, with feeble pulse, a
smaller dose should be given. RINGER.