

SCIATICA (Continued).

1236—R Pulv. sulphuris sublimati, . . . ℥iv.

Sig.: Dust thickly on the limb and envelop it in soft flannel.

RINGER.

1237—R Veratriæ, ℥i-ij.
Adipis, ℥j.—M.

Ft. unguentum.

Sig.: Rub into the painful part for half an hour, twice daily.

TURNBULL.

1238—R Emplastri cantharidis, 1 in. by 5 in.

Sig.: Apply over the course of the nerve, until it is blistered; then apply poultices until the blister is well filled; then cut and dress with—

1239—R Morphiæ acetatis, gr. ij.
Pulv. acaciæ, gr. x.—M.

Sig.: Use locally.

HARTSHORNE.

1240—R Antipyrin., ℥ij.
Syr. aurantii cort., ℥ss.
Aquæ aurantii flor., ad ℥ij.—M.

Sig.: A dessertspoonful every hour to four hours, until three to six doses are taken.

GERMAIN SÉE, Annual Univ. Med. Sci.

1241—R Methyl chloridi, ℥ss.

Sig.: Apply with an atomizer, locally, but with care.

DEBOVE, Annual Univ. Med. Sci.

1242—R Chloroformi, ℥j.
Olei vaselini, ℥iv.—M.

Sig.: Inject fifteen to twenty minims hypodermically, and repeat if necessary.

MEUNIER, Annual Univ. Med. Sci.

1243—R Saloli, ℥ss.
Olei vaselini, ℥v.—M.

Sig.: Inject twenty or thirty minims hypodermically over the course of the nerve.

MEUNIER, Annual Univ. Med. Sci.

1244—R Saloli,
Sacchari lactis, āā ℥iij.—M.

In pulv. no. xii div.

Sig.: A powder every four to six hours.

ASCHENBACH, Annual Univ. Med. Sci.