

SEA-SICKNESS (Continued).

1259—R Chloral hydratis, ℥ss.
Syr. aurantii cort., ℥j.
Aquæ aurantii flor., ad ℥ij.—M.

Sig.: One or two teaspoonfuls in water every four hours. RINGER.

1260—R Chloroformi, ℥ss.

Sig.: Two to five minims on sugar every half-hour until relieved. BARTHOLOW.

1261—R Sodii bromidi, ℥j.
Ammonii bromidi, ℥ss.
Aquæ menthæ pip., ℥v.—M.

Sig.: A tablespoonful before meals and at bed-time. To be used for three days before embarking.

BEDARD.

1262—R Cocaini hydrochloratis, gr. xxx.
Aquæ destillatæ, ℥ivss.—M.

Sig.: Four or five drops on a small piece of ice thrice daily. W. OTTO.

1263—R Antipyrin., gr. lxxv.
Cocaini hydrochloratis, gr. iss.
Caffeini, gr. iv.
Strychniæ sulphatis, gr. $\frac{1}{30}$.
Spts. vini gallici, ℥iiss.
Aquæ destillatæ, ℥xxiiss.—M.

Sig.: A tablespoonful before embarking, and two others during the day, or three during the twenty-four hours. ROUQUETTE.

SEPTICÆMIA (See Pyæmia).

SHINGLES (See also Skin Diseases and Herpes Zoster).

1264—R Zinci phosphidi,
Ext. nucis vomicæ, āā gr. x.—M.

Ft. massa et in pil. no. xxx div.

Sig.: One pill every two to four hours. BULKLEY.

1265—R Pulv. amyli, ℥iv.

Sig.: Dust over the eruption and on a muslin band sewed tightly around the body, to protect it from the friction of the clothes. BULKLEY.