

**SKIN DISEASES (Continued).**

1304—R Pulv. camphoræ, . . . . . ℥ss.  
 Pulv. zinci oxidi, . . . . . ℥ij.  
 Glycerinæ, . . . . . ℥xl.  
 Unguenti benzoini, . . . . . ℥j. —M.

Sig.: Apply locally at once, or some isolating powder, as talc, bismuth, or lycopodium may be used first. (*In vesiculous eczema.*) DUHRING.

1305—R Liq. potassii arsenitis, . . . . . ℥j.  
 Vini ferri amari, . . . . . ℥iss.  
 Syr. tolutani, . . . . . ℥ij.  
 Aquæ anethi, . . . . . ℥ij.—M.

Sig.: A teaspoonful or two thrice daily. (*In eczema.*) ERASMUS WILSON.

1306—R Ungt. zinci oxidi,  
 Ungt. plumbi subacetatis, āā ℥ss.  
 Chloral hydratis,  
 Pulv. camphoræ, . . . . . āā gr. xv.—M.

Ft. ungt.

Sig.: Use two or three times daily, after bathing with warm water. (*In general eczema.*) GROSS.

1307—R Olei morrhuæ, . . . . . ℥ij.  
 Vitelli ovi, . . . . . no. j.  
 Liq. sodii arseniatis, . . . . . ℥j.  
 Syrupi simplicis, . . . . . ℥ij.  
 Aquæ, . . . . . ℥iv.—M.

Sig.: A half-teaspoonful thrice daily. (*For chronic infantile eczema.*) DOYON.

1308—R Ferri et ammonii citratis, . . . . . ℥j.  
 Potassii citratis, . . . . . ℥ij.  
 Liq. potassii arsenitis, . . . . . ℥i-ij.  
 Tinct. nucis vomicæ, . . . . . ℥ij.  
 Tinct. cinchonæ co., . . . . . ad ℥iv.—M.

Sig.: A teaspoonful in water after meals, as a tonic and alterative. (*In eczema.*) BULKLEY.

1309—R Resorcin.,  
 Zinci oxidi, . . . . . āā ℥j.  
 Ungt. aquæ rosæ, . . . . . ℥x.—M.

Ft. ungt.

Sig.: Apply locally. (*In chronic indurated eczema of infants.*) FLIESBURG, Annual Univ. Med. Sci.