

SKIN DISEASES (Continued).

1366—℞ Potassii cyanidi, gr. iv.
 Chloroformi, ℥viiij.
 Glycerinæ, ℥j.
 Cerati simplicis, ℥vj.—M.

Ft. ungt.
 Sig.: Apply locally. (*In lichen agrius. Also in pruritus.*)
 NELIGAN.

1367—℞ Liq. plumbi subacetatis, ℥i-iiij.
 Infusi althææ, Oj. —M.

Ft. lotio.
 Sig.: Apply locally. (*In lichen agrius. Also in chronic eczema.*)
 BURGESS.

1368—℞ Bismuthi subnitratiss, ℥ij.
 Pulv. zinci oxidi, ℥ij.
 Tinct. digitalis, ℥ss.
 Aquæ, q. s. ad ℥vj.—M.

Ft. lotio.
 Sig.: To be used as a lotion, after an alkaline bath, as bicarbonate of soda with bran. (*In lichen planus.*)
 TILBURY FOX.

1369—℞ Sodii carbonatis, ℥j.
 Aquæ rosæ, ℥vj.
 Glycerinæ, ℥ij.—M.

Ft. lotio.
 Sig.: Apply to the eruption. (*In infantile lichen or strophulus.*)
 TILBURY FOX.

1370—℞ Pulv. lycopodii, ℥ss.

Sig.: Use as a dusting powder after the bullæ are cut. Then use zinc ointment or an astringent solution; and then use the following:—

1371—℞ Argenti nitratis, gr. iii-iv.
 Adipis, ℥j. —M.

Ft. ungt.
 Sig.: Apply locally. (*In chronic ulceration following pemphigus.*)
 TILBURY FOX.

1372—℞ Unguenti hydrargyri nitratis, ℥ij.
 Unguenti simplicis, ℥vj.—M.

Sig.: Use twice daily. Also take internally five drops of Fowler's Solution, well diluted, thrice daily after meals. (*In pemphigus.*)
 WARING.