

SKIN DISEASES (Continued).

1417—℞ Acidi carbolici, ℥j.
Glycerinæ, ℥ss-j.—M.

Ft. lotio.

Sig.: Use locally night and morning, rubbing in well. (*In tinea tonsurans.*) TILBURY FOX.

1418—℞ Sulphuris loti, ℥ij.
Spts. camphoræ, ℥ss.
Glycerinæ, ℥ss.
Hydrargyri bisulphidi, ℥ss.
Pulv. amyli, ℥ij.
Aquæ, q. s. ad Oj.—M.

Ft. lotio.

Sig.: Use locally night and morning. (*In tinea tonsurans.*) STARTIN.

1419—℞ Hydrargyri ammoniati,
Hydrargyri oxidi rubri, āā gr. vj.
Adipis, ℥j. —M.

Ft. ungt.

Sig.: Use after epilation and washing. (*In tinea tonsurans.*) STARTIN.

1420—℞ Acidi acetici glacialis, ℥j.
Sig.: Use as a paint, once or more, and dry off with blotting-paper if it produces much irritation. (*In tinea circinata.*) TILBURY FOX.

1421—℞ Hydrargyri nitratis, ℥iss.
Adipis, ℥j.—M.

Ft. ungt.

Sig.: Rub in, night and morning, for a day or two. (*In tinea circinata.*) TILBURY FOX.

1422—℞ Hydrargyri chloridi corr., gr. ij.
Adipis, ℥j.—M.

Ft. ungt.

Sig.: Use locally, rubbing in well. (*In tinea circinata.*) TILBURY FOX.

1423—℞ Acidi sulphurosi, ℥ij.
Aquæ, ℥viiij.—M.

Ft. lotio.

Sig.: Wash the surface with soap and water, and apply the lotion, night and morning, on compresses of lint covered with oiled silk for at least one hour. (*In tinea circinata, where the disease is more or less general.*) TILBURY FOX.