

SKIN DISEASES (Continued).

1431—R Sodii hyposulphitis, . . . ℥iv-vj.
Aquæ, . . . ℥vj. —M.

Ft. lotio.

Sig.: Wash the parts well with yellow soap (sapo terebinthinæ), then sponge with weak vinegar and water, and apply the lotion freely, even after the disease has vanished. (*In tinea versicolor.*)

TILBURY FOX.

1432—R Acidi salicylici, . . . gr. xxx.
Sulphuris loti, . . . ℥iiss.
Lanolini, . . . ℥xxv.—M.

Ft. ungt.

Sig.: Apply with friction. (*In tinea versicolor.*)

LIEBREICH.

1433—R Acidi salicylici, . . . gr. xlv.
Sulphuris loti, . . . ℥iiss.
Lanolini,
Vaselini, . . . āā ℥xiiiss.—M.

Ft. ungt.

Sig.: After rubbing well with tar-soap, rub in the ointment well every evening. Wash off in the morning. (*In tinea versicolor.*)

E. BESNIER.

1434—R Acidi carbolic, . . . ℥j.
Glycerinæ, . . . ℥ix.—M.

Ft. sol.

Sig.: Cut the eyelashes short. Scrape off all crusts and the surface of the exposed ulcerations. Paint the raw surface with the solution. Dress with iodoform. Remove the dressing in two or three days. (*In tinea tarsi.*)

TEALE, Annual Univ. Med. Sci.

1435—R Sodii bicarbonatis, . . . ℥ii-x.
Aquæ ferventis (90°-95° F.),
cong. xx-xxx.—M.

Sig.: Alkaline bath. (*In eczema, psoriasis, urticaria, lichen, and prurigo, where there is much local irritation.*)

TILBURY FOX.

1436—R Potassii carbonatis, . . . ℥ii-vj.
Sodii boratis, . . . ℥iij.
Aquæ ferventis (90°-95° F.),
cong. xx-xxx.—M.

Sig.: Alkaline bath. Use the same as the preceding.

TILBURY FOX.