

SMALL-POX (Continued).

1444—R Collodii flexilis, ℥j.
Sig.: Apply every day or two with a brush to the eruption. (*To prevent pitting.*) RINGER.

1445—R Liq. ammonii acetatis, ℥iiss.
Spts. ætheris nitrosi, ℥ss.—M.
Sig.: A tablespoonful every two or three hours in a wineglassful of water. When the fever abates give one or two grains of quinine every three or four hours, with a one-grain opium pill at night. HARTSHORNE.

1446—R Acidi carbolici,
Acidi acetici, āā ℥i—iss.
Tinct. opii,
Spts. chloroformi, āā ℥j.
Aquæ destillatæ, ad ℥viiij.—M.
Sig.: A tablespoonful every four hours until the fever subsides. NAPHEYS.

1447—R Calcis calcinatæ, ℥iv.
Sulphuris, ℥viiij.
Aquæ, Ov.—M.
[Boil in an earthenware pan, evaporate to three pints, and filter.]
Sig.: Apply locally as a lotion to the eruption. (*To prevent pitting and secondary fever.*) PETERS.

1448—R Sodii salicylatis, ℥ij.
Glycerinæ, ℥j.
Aquæ menthæ pip., ad ℥iiij.—M.
Sig.: One or two teaspoonfuls three or four times daily. (*Aborts the pustules.*) REIMER.

1449—R Hydrargyri chloridi corrosiv., gr. ii—iv.
Aquæ, ℥vj. —M.
Ft. lotio.
Sig.: Wet compresses and apply to the eruption. SKODA.

SPERMATORRHŒA.

1450—R Ext. belladonnæ,
Pulv. belladonnæ, āā gr. iij.
Confect. rosæ, q. s. ut ft. massa.—M.
In pulv. nō. x div.
Sig.: From one to three pills at bed-time, and during the day from fifteen to sixty grains of potassium bromide in divided doses. (*When due to spasm of the vesiculæ seminales.*) SINÉTY.