

STRANGURY (Continued).

1466—R Ext. belladonnæ, . . . gr. i-iv.
Olei theobromæ, . . . ℥ss. —M.

Ft. suppositorium no. i.

Sig.: Introduce into the bowel, and repeat in four hours, if it be necessary. HARTSHORNE.

STRUMA (See Rachitis).

SUPPURATION (See Abscess).

SWEATING (See Phthisis and Fetor).

SYCOSIS (See Tinea in Skin Diseases).

SYNOVITIS.

1467—R Iodini, . . . ℥iv.
Potassii iodidi, . . . ℥j.
Aquæ destillatæ, . . . ℥vj.—M.

Sig.: Apply externally, with a brush. LUGOL.

1468—R Ungt. hydrargyri, . . . ℥ij.
Pulv. ammonii chloridi, . . . ℥j.—M.

Sig.: For inunction. DUPUYTREN.

1469—R Morphiæ, . . . gr. viij.
Hydrarg. oleatis (5 per cent.
to 10 per cent.), . . . ℥j.—M.

Sig.: Apply twice daily with a soft brush. (*In the acute form.*) MARSHALL.

1470—R Empl. cantharidis, . . . 1 in. by 2 in.

Sig.: Apply every night until the skin is well reddened. If this does not avail, leave on until a bleb is formed, which may be cut, poulticed, and dressed with simple cerate. (*In the chronic form.*) RINGER.

1471—R Saponis mollis, . . . ℥ij.
Alcoholis vel aquæ cologn., . . . ℥j.—M.

Sig.: Soak linen rags in the solution and apply about the joint. KAPPESSER.

1472—R Vitelli ovi, . . . no. ij.
Pulv. sacchari albi, . . . ℥iv.
Olei amygdalæ amaræ, . . . gtt. ij.
Aquæ aurantii flor., . . . ℥ij.
Olei morrhuæ, . . . ℥v.—M.

Sig.: From a teaspoonful to a tablespoonful thrice daily. (*In strumous synovitis.*) HEDER.