

SYNOVITIS (Continued).

1473—R Tinct. iodinii, ℥j.
Sig.: Apply with a brush every second or third day.
RINGER.

STYE (Hordeolum).

1474—R Acidi boracici, ℥iv.
Aquæ destillatæ, ℥v.—M.
Ft. lotio.
Sig.: Apply to the eyelids several times daily.
ABADIE, Annual Univ. Med. Sci.

SYPHILIS.

1475—R Hydrargyri salicylatis, gr. viiss.
Confectionis rosæ, ℥ss. —M.
Ft. massa et in pil. no. lx div.
Sig.: One thrice daily, after meals.
CHAVES, Annual Univ. Med. Sci.

1476—R Hydrargyri carbolatis, gr. xvij.
Ext. glycyrrhizæ,
Pulv. glycyrrhizæ, āā q. s. ut ft. massa.—M.
Ft. massa et in pil. no. lx div.
Obduc. bals. tolutano.
Sig.: Two to four pills daily.
SZADEK, Annual Univ. Med. Sci.

1477—R Hydrargyri chloridi mitis, gr. xij.
Olei vaselini, ℥ccxxv.
—M.
Sig.: For hypodermic injection. Twenty to thirty
minims to be used.
BALZER, Annual Univ. Med. Sci.

1478—R Hydrargyri chloridi mitis,
Sodii chloratis, āā gr. xv.
Aquæ destillatæ, ℥iiss.—M.
Sig.: For hypodermic injection. Twenty to thirty
minims to be used.
KRECKE.

1479—R Hydrargyri chloridi mitis, gr. xv.
Olei olivæ, ℥iiss.—M.
Sig.: For hypodermic injection. Twenty to thirty
minims to be used.
KOPP.