

SYPHILIS (Continued).

1480—R Hydrargyri chloridi mitis, . gr. iss.
Glycerinæ, ℥xv.—M.

Sig.: For hypodermic injection. To be used each time. Two or three injections required for an average case, before the symptoms yield. SCARENZIO.

1481—R Hydrargyri oxidi flavæ, . ʒiss.
Hydrargyri chloridi corros., . gr. $\frac{3}{4}$.
Glycerinæ puræ, ʒxxv.—M.

Sig.: For hypodermic use. Twelve and a half minims to be used for each injection. DE SMET.

1482—R Hydrargyri chloridi corros.,
Ammonii chloridi, āā gr. iij.
Aquæ destillatæ, ʒiss.—M.

Ft. sol et adde—

Albuminis ovi, ʒiss.
Aquæ destillatæ, ʒv.

Misce, cola, et adde—

Aquæ destillatæ, q. s. ad ft. ʒx.—M.

Sig.: For hypodermic use. Three to ten minims to be used for each injection. ℥j contains corrosive sublimate, gr. $\frac{1}{200}$. POTTER.

1483—R Hydrargyri protoiodidi,
Lactucarii, āā gr. xv.
Ext. opii, gr. ii $\frac{1}{4}$.
Ext. guaiaci, ʒss. —M.

Ft. massa et in pil. no. xx div.

Sig.: One pill at breakfast and after supper, followed by a draught of water. DIDAY.

1484—R Hydrargyri chloridi corros.,
Ammonii chloridi, āā gr. iiss.
Aquæ destillatæ, ʒiv. —M.

Ft. sol. et adde—

Potassii iodidi, ℥j.
Aquæ destillatæ, ʒxvj.—M.

Sig.: A tablespoonful before each meal. BESNIER.

1485—R Pil. hydrargyri, ʒij.
Ferri sulphatis exsiccatae, ʒj.
Ext. opii aquosi, gr. v.—M.

In pil. no. xx div.

Sig.: One pill thrice daily. F. N. OTIS.