

TRISMUS NEONATORUM (See also *Tetanus*).

1528—R Tinct. opii, ℥j.
Olei ricini, ℥j.—M.

Sig.: A teaspoonful every four hours, with a warm bath. DRUITT.

1529—R Tinct. opii, ℥iv.

Sig.: One drop every three hours, alternating with the following:—

1530—R Pulv. ipecac. co., gr. xxiv.
Zinci sulphatis, ℥j.—M.

In pulv. no. xii div.
Sig.: A powder every three hours. FURLONGE.

1531—R Tinct. opii, gtt. v.
Tinct. asafœtidæ, ℥iss.
Syr. simplicis, ℥v.
Aquæ, ad ℥xv.—M.

Sig.: A half-teaspoonful hourly. EBERLE.

1532—R Chloral hydratis, gr. i-ij.
Syr. simplicis, ℥j.—M.

Sig.: To be given every two hours, unless there be profound sleep. Double the dose, if given by the rectum. WIDERHOFER.

1533—R Ext. gelsemii fld., ℥viii-xvj.
Syr. simplicis, ℥j.
Aquæ, q. s. ad ℥iv.—M.

Sig.: A half-teaspoonful every two to four hours. BARTHOLOW.

TUBERCULOSIS (See *Rachitis* and *Phthisis*).

TYMPANITES (See also *Fever*).

1534—R Pulv. capsici, gr. vi-xxiv.
Sacchari lactis, ℥iss.—M.

In pulv. no. xii div.
Sig.: A powder every four hours. PHILLIPS.

1535—R Olei terebinthinæ, ℥j.
Pulv. acaciæ, q. s.
Misce, et adde—
Decocti hordei, ℥xix.—M.

Ft. enema.
Sig.: Inject into the bowel. HOOPER.