

**WHOOPING-COUGH.**

1612—R Acidi carbolici, . . . . . ℥ss.  
Potassii chloratis, . . . . . ℥ij.  
Glycerinæ, . . . . . ℥iv.  
Aquæ, . . . . . q. s. ad ℥vj.—M.

Sig.: Use with a steam atomizer three times daily.  
J. LEWIS SMITH

1613—R Sodii benzoatis, . . . . . ℥iv.  
Aquæ menthæ pip., . . . . .  
Aquæ destillatæ, . . . . . āā ℥x.  
Syr. aurantii, . . . . . ℥ij.—M.

Sig.: A dessertspoonful every hour or two.  
LETZERICH

1614—R Ext. aconiti, . . . . . gr. j.  
Syr. ipecac., . . . . . ℥xliv.  
Aquæ lauro-cerasi, . . . . . ℥j.  
Mucilag. acaciæ, . . . . . ℥viss.—M.

Sig.: A teaspoonful to a tablespoonful, according to  
age, every hour. DERVIEUX

1615—R Pulv. acidi boracici, . . . . . ℥i-iss.  
In pulv. no. xxx div.

Sig.: Insufflate one powder into each nostril, three  
times during the day and once at night.  
HOLLOWAY, Annual Univ. Med. Sci.

1616—R Quiniæ sulphatis, . . . . . gr. l.  
Acidi sulphurici, . . . . . gtt. xxx.  
Aquæ destillatæ, . . . . . ℥v ℥v.—M.

Sig.: Use as a spray to the fauces every two hours  
for the first three days, and every three hours for the  
remainder of the first week, after which it will be  
unnecessary. KOLOVER, Annual Univ. Med. Sci.

1617—R Pulv. benzoini,  
Bismuthi salicylatis, . . . . . āā ℥iiss.  
Quiniæ sulphatis, . . . . . gr. xxx.—M.

Sig.: Insufflate the fauces several times daily.  
MOIZARD, Annual Univ. Med. Sci.

1618—R Antipyrin., . . . . . gr. ij.  
Sacchari albi, . . . . . ℥j.—M.

In pulv. no. xiv div.  
Sig.: A powder three times during the day and once  
during the night, for very young children. Dose  
increased up to fifteen grains for adults.  
SONNENBERGER, Annual Univ. Med. Sci.