

WORMS (Continued).

1626—R Pelletierine sulphatis, . . . gr. vi-viiss.
Pulv. acidi tannici, . . . gr. viiss.
Syr. simplicis, . . . ℥ij. —M.

Sig.: Take a little milk for supper, and a simple enema at bed-time. Take the mixture the following morning before breakfast, and lie down to prevent vertigo. In a quarter- or a half-hour take an ounce of castor oil. (*For tape-worm.*) LABBÉ.

1627—R Thymoli, . . . ℥ij.

In pulv. no. xii div.

Sig.: A powder every fifteen minutes. A dose of castor oil should precede and follow the powders. (*For tape-worm.*) N. CAMPI, Annual Univ. Med. Sci.

1628—R Pelletierine tannatis, . . . gr. vij.
Syr. simplicis, . . . ℥ss. —M.

Sig.: Milk diet the day before. Before breakfast, a dose of infusion of senna. One hour later, half the medicine, and the rest a half-hour later. A half-hour later an ounce of castor oil. Patient to remain in bed during treatment. A large vessel of warm water should be ready to receive the worm. (*For tape-worm.*) BÉRENGER FÉRAUD.

1629—R Chloroformi, . . . ℥j.
Syr. simplicis, . . . ℥j.—M.

Sig.: To be given in three doses, at intervals of two hours; taken fasting, and followed by castor oil. (*For tape-worm.*) THOMPSON, Annual Univ. Med. Sci.

1630—R Oleoresinæ filicis maris,
Tinct. vanillæ, . . . āā ℥xliv.
Syr. terebinthinæ, . . . ℥vi ℥xv.
Pulv. acaciæ, . . . gr. xxx.
Aquæ destillatæ, . . . ℥vj ℥xv.—M.

Sig.: To be taken at one dose, in an equal quantity of milk. Castor oil should be given a few hours later. (*For tape-worm.*)

EILLARD, Annual Univ. Med. Sci.

1631—R Pulv. kamalæ, . . . gr. v-x.
Syr. aurantii, . . . ℥ss.
Mucil. tragacanthæ, . . . ℥j.
Aquæ, . . . ℥j.—M.

Ft. haustus.

Sig.: Take early in the morning, and four hours later a purge. For a child of two to five years. (*In tape-worm.*) T. H. TANNER.