

WORMS (Continued).

1632—℞ Olei terebinthinæ,
Oleoresinæ filicis maris, āā ʒj.
Mucilag. acaciæ, ʒij.—M.

Ft. emulsio.

Sig.: Day before treatment, a milk or thin soup diet, and one drachm of compound jalap-powder. The emulsion is taken the following morning, fasting, and a half-hour later a dose of castor oil. (*For tape-worm.*)

F. A. A. SMITH, Annual Univ. Med. Sci.

1633—℞ Tinct. rhei, gtt. iij.
Tinct. zingiberis, gtt. j.
Magnesii carbonatis, ʒv.
Aquæ, ʒij.—M.

Sig.: This dose should be taken three or four times daily, according to the effect on the bowels. (*For oxyuris.*)

SYDNEY MARTIN, Annual Univ. Med. Sci.

1634—℞ Sodii chloridi, ʒx.
Aquæ, ʒvj.—M.

Ft. sol.

Sig.: To be injected by the rectum. (*For oxyuris.*)

EILLARD, Annual Univ. Med. Sci.

1635—℞ Acidi tannici, gr. xv.
Olei theobromæ, ʒj. —M.

Ft. suppositorium no. i.

Sig.: To be introduced into the rectum. (*For thread-worms.*)

EILLARD, Annual Univ. Med. Sci.

1636—℞ Tinct. ferri chloridi, ʒss.
Aquæ, ʒj.—M.

Sig.: One-fourth to one-third, as a rectal enema. (*For seat-worms.*)

RINGER.

1637—℞ Ext. spigeliæ et sennæ fld., ʒj.
Santonini, gr. viij.—M.

Sig.: A teaspoonful to a child of five years. (*For ascarides and lumbrici.*)

J. LEWIS SMITH.

1638—℞ Trochisci santonini (U. S. P), no. xxiv.

Sig.: One to six at bed-time for children, with a dose of castor oil the following morning. (*For ascarides.*)

BARTHOLOW.

1639—℞ Infusi quassiaæ, ʒvj.

Sig.: Use as a rectal injection. (*For seatworms.*)

RINGER.