

WORMS (Continued).

1640—R Santonini, gr. xij.
Olei theobromæ, ℥j. —M.

Fiant suppositoria no. iv.

Sig.: One at bed-time, introduced into the bowel.

(For seat-worms.)

HARTSHORNE.

1641—R Olei chenopodii, gtt. lx-℥j.
Mucilag. acaciæ, ℥ij.
Syr. simplicis, ℥j.
Aquæ cinnamomi, ℥ij.—M.

Sig.: A dessertspoonful thrice daily for three days,
and repeat after three days. For a child of two years.

(For lumbrici.)

MEIGS and PEPPER.

1642—R Hydrargyri chloridi mitis, gr. ij.
Santonini, gr. iss.
Sacchari lactis, gr. xv.—M.

Sig.: To be taken in honey at one dose, by a child
of two years. (For stomach-worms.)

E. BOUCHUT.

WOUNDS.

1643—R Acidi tannici, ℥iiss.
Alcoholis absoluti, ℥ss.
Ætheris, ℥iiss.
Collodii, ℥xij.—M.

Sig.: "Styptic colloid."

1644—R Acidi carbolici, ℥ss.
Olei ricini, āā ℥ss.
Collodii, ℥j.—M.

Sig.: "Carbolized collodion."

1645—R Hydrargyri chloridi corrosivi, gr. viiss.
Aquæ ferventis, Oij. —M.

Sig.: "Sublimate solution (1 to 2000)." (For wash-
ing wounds, irrigating cavities, or saturating dressings.)

1646—R Pulv. acidi salicylici, ℥j.

Sig.: Use as a dusting powder.

THIERSCH.

1647—R Iodoformi, ℥j.
Collodii flexilis, ℥vij.—M.

Sig.: Hold or stitch the edges of the wound together,
and apply with a brush. (In superficial wounds.)

BRUNS, Annual Univ. Med. Sci.