

WOUNDS (Continued).

1648—R Iodoformi, . . . . . gr. c.  
Thymoli, . . . . . gr. cc.  
Sacch. lactis, . . . . . gr. j.—M.

Ft. pulv.

Sig.: Apply as a powder thrice daily.

1649—R Ext. calendulæ fld., . . . . . ℥j.  
Alcoholis, . . . . . ℥ij.—M.

Sig.: Use as a dressing.

R. G. REYNOLDS, Annual Univ. Med. Sci.

1650—R Iodol, . . . . . ℥j.  
Glycerinæ, . . . . . ℥j.  
Vaselini, . . . . . ℥vij.—M.

Ft. ungt.

Sig.: Use locally.

WOLFENDEN, Annual Univ. Med. Sci.

1651—R Pulv. naphthol., . . . . . ℥j.

Sig.: Use as a dusting powder.

BOUCHARD, Annual Univ. Med. Sci.

1652—R Acidi boracici, . . . . . ℥iiss.  
Ess. eucalypti, . . . . . ℥iiss.  
Vaselini, . . . . . ℥xxv.—M.

Ft. ungt.

Sig.: To be used as a dressing. (*The boracic acid may be replaced by mercuric chloride corr., gr. iss, if desired.*)

BRONDEL.

1653—R Pulv. saloli, . . . . . āā ℥ss.—M.  
Pulv. amyli, . . . . . āā

Sig.: Use as a dusting powder.

CREYX et JARRY.

1654—R Saloli, . . . . . āā ℥j.  
Ætheris, . . . . . āā  
Solve et adde—  
Collodii flexilis, . . . . . ℥viiss.—M.

Sig.: Use with a camel's hair brush.

NICOT.

1655—R Acidi boracici, . . . . . ℥iiss.  
Glycerinæ, . . . . . ℥ij.  
Infusi caryophylli, . . . . . Oij.  
(℥iv ad Oj.)  
Olei menthæ pip., . . . . . ℥vj.—M.

Sig.: "Aseptin."

MAGNUS TROILIUS.