

minims of the weaker, and gradually increasing, is recommended. Chrysophanic acid, one-fifteenth to one-seventh grain, is highly recommended in eczema, lichen, prurigo, and psoriasis. Pilocarpin hypodermically is also of use.

*Enuresis, Nocturnal.*—Two very small doses of the nitrate of strychnia, injected in the vicinity of the rectum at suitable intervals, have proved successful.

*Epilepsy.*—Curare, in solution, seven grains in seventy-five minims of water, with two drops of hydrochloric acid. About once a week inject eight drops beneath the skin. It has cured cases of several years' standing within two months. Also lobeline hydrobromate, one hundredth to one-twentieth grain, hypodermically.

*Erysipelas.*—Carbolic acid, three per cent. solution, eight or ten injections at the same time, so as to surround and cover the inflamed regions. Also, salicylic acid in the same manner.

*Fevers.*—Antipyrine hypodermically is very valuable as an antipyretic in doses of ten or fifteen grains. Cocaine, one-eighth grain, may be added. Thallin sulphate, one to one and a half grains, also useful.

*Fractures Ununited.*—Glacial acetic acid, five to ten minims between the ends of the bones with hypodermic syringe. Iodine has also succeeded, used in the same way.

*Foreign Body in Oesophagus.*—Threatened strangulation from impaction of the gullet has been promptly relieved by inducing vomiting; apomorphia, one-tenth grain, hypodermically. Emetia is also suggested in the same way.

*Goitre* has been successfully treated by subcutaneous injections of ergotine, one-third grain, gradually increased to one grain.

*Hæmoptysis.*—Sclerotinic acid, as a substitute for ergotine, five per cent. solution, injected in the neck or arm.

*Heart Failure.*—Tincture of digitalis, ten to thirty minims. Digitaline, and helleborein (Mercks) have been of great use. (See formulæ.) Brandy and whisky also used hypodermically.

*Hemorrhages.*—Hæmoptysis, hæmatemesis, and uterine hemorrhages have all been arrested by hypodermics of ergotine or sclerotic acid. In pain, add morphia.

*Hemorrhoids.*—Iodine, carbolic acid, perchlor. iron, and other preparations have been used successfully—a few drops of either injected into each pile—usually operating on only one at a session, waiting several days before repeating. For details see *The Monthly Review of Medicine and Pharmacy*, July, 1879.

*Hernia* is more easily reduced by giving a hypodermic of morphia with or without atropia.

*Hiccough.*—In an obstinate case, resisting all other means, three-eighths of a grain of hydrochlorate of pilocarpin, hypodermically, quickly proved successful.

*Hydrophobia.*—Much amelioration of the symptoms has followed hypodermics of curare.

*Mania and Melancholia.*—Paraldehyde, six to twelve minims in solution, highly commended. Hyoscyamia, one ninety-sixth grain, and hyoscine hydrobromate, one two-hundredth to one-ninetieth grain, also very useful.

*Nasal Polypus.*—Carbolic acid, one part, glycerine, four parts; twenty drops injected into the tumor effectually dissipated the polypus in case reported.

*Neuralgias.*—Osmic acid, one-fifteenth to one-seventh grain, is recommended by Billroth and Neuber for obstinate cases, also for interstitial injection of tumors. Also chloroform and antipyrine, hypodermically.

*Night-Sweats.*—Atropia has given good results in injections of about one-fortieth of a grain at bed-time. Agaricin, one-tenth grain, has proved very effectual.

*Opium Habit.*—Sparteine sulphate has been used hypodermically.

*Paralysis.*—Strychnia, used with great success hypodermically.

*Peritonitis.*—Morphia and conia have been of service hypodermically.